Sobering Facts: Alcohol-Impaired Driving



KEEP IOWA SAFE

Keep alcohol-impaired drivers off the road.


This fact sheet provides a snapshot of **alcohol-impaired driving deaths** and an overview of proven strategies to reduce or prevent alcohol-impaired driving. The information can help decision makers and community partners see gaps and identify relevant strategies to address the problem of alcohol-impaired driving.

FAST FACTS

- Drivers with a blood alcohol concentration (BAC) above the state's legal limit are considered alcohol-impaired by law.
- More than 10,000 people in the United States die each year in crashes that involve an alcohol-impaired driver.
- Because of dedicated efforts, rates of alcohol-impaired driving and alcohol-impaired fatal crashes in the United States have gone down since the 1980s.
- Still, alcohol-impaired drivers get behind the wheel millions of times each year.

ALCOHOL-IMPAIRED DRIVING DEATH RATES BY AGE

Per 100,000 Population



Deaths in crashes involving a driver with BAC ≥ 0.08%. Source: Fatality Analysis Reporting System (FARS), 2018

★ Fatality rates based on fewer than 20 deaths are suppressed.

ALCOHOL-IMPAIRED DRIVING DEATH RATES BY SEX

Per 100,000 Population



Deaths in crashes involving a driver with BAC $\geq 0.08\%$. Source: Fatality Analysis Reporting System (FARS), 2018

PERCENTAGE OF ADULTS WHO REPORT DRIVING AFTER DRINKING TOO MUCH

In the Past 30 Days

1.7%

3.0%

Source: Behavioral Risk Factor Surveillance System (BRFSS), 2018



WHAT WORKS

The strategies in this section are effective for reducing or preventing alcohol-impaired driving. They are recommended by *The Guide to Community Preventive Services* and/or have been demonstrated to be effective in reviews by the National Highway Traffic Safety Administration.* Different strategies might require different resources for implementation or have different levels of impact. **Find strategies that are right for your state.**

Strategies to Reduce or Prevent Alcohol-Impaired Driving

- Alcohol-impaired driving laws make it illegal to drive with a BAC at or above
 a specified level (0.05% or 0.08%, depending on the state). For people under
 21, zero tolerance laws make it illegal to drive with any measurable amount of
 alcohol in their systems. These laws, along with laws that maintain the minimum
 legal drinking age at 21, are in place in all 50 states and DC and have saved tens
 of thousands of lives.
- Publicized sobriety checkpoints allow police to briefly stop vehicles at specific, highly visible locations to check drivers for impairment. Police may stop all or a certain portion of drivers. Sobriety checkpoints should be well publicized (e.g., through mass media campaigns) and conducted regularly for greatest impact.
- High-visibility saturation patrols consist of a large number of police patrolling
 a specific area, usually at times and locations where alcohol-impaired driving
 crashes are more common. Like sobriety checkpoints, these patrols should be
 well publicized and conducted regularly.
- Ignition interlocks for all, including first-time, convicted offenders can be
 installed in vehicles to measure alcohol on drivers' breath. Interlocks keep vehicles
 from starting if drivers have a BAC above a certain level, usually 0.02%. Interlocks
 are highly effective at preventing repeat offenses while installed. Incorporating
 alcohol problem assessment and treatment into interlock programs shows
 promise in reducing repeat offenses once interlocks are removed.
- Alcohol problem assessment and treatment programs can be used for those
 arrested for alcohol-impaired driving. Treatment is most effective when combined
 with other sanctions and when offenders are closely monitored. Assessment and
 treatment are critical to the success of **DWI courts**, which are specialized courts
 focused on changing the behavior of alcohol-impaired driving offenders.
- Alcohol screening and brief interventions take advantage of "teachable moments" and can be delivered in health care, university, and other settings to identify people at risk for alcohol problems and get them treatment as needed.
- Multi-component interventions combine several programs or policies to
 prevent alcohol-impaired driving. The key to these comprehensive efforts is
 community mobilization, which involves coalitions or task forces in design and
 implementation.
- School-based instructional programs are effective at teaching teens not to ride with alcohol-impaired drivers.

IN IOWA:

- It is illegal to drive with a BAC at or above 0.08%.
- Publicized sobriety checkpoints are not allowed.
- Ignition interlocks are required for all (including first-time) convicted offenders.

For up-to-date information on laws in your state, check with the Insurance Institute for Highway Safety at www.iihs.org.



Find more information at www.cdc.gov/motorvehiclesafety

- Injuries, costs, and other data related to alcohol-impaired driving.
- Detailed information on effective strategies to reduce or prevent alcohol-impaired driving.
- An interactive calculator to estimate the expected number and monetized value of injuries prevented, lives saved, and costs of implementation for 14 effective interventions.